



Scheme 16.2



The Cu^{II} -bound alkoxide 13 then reacts with 7 to afford the aldehyde 6 (Step III). The mechanism of this process is not known, but it could occur via a homolytic mechanism as shown in Equation 16.23, where the cupric center is reduced to cuprous ion, resulting in the formation of 10 and 12. Overall, the conversion of the alcohol to the aldehyde is a net two-electron oxidation reaction, with the cupric atom and TEMPO (7) serving as complementary one-electron oxidants.



In the experiment described in this section, a substituted benzylic alcohol 5 will be oxidized under aerobic conditions to the corresponding benzaldehyde 6 using the Cu/TEMPO catalyst system. For additional information, you may wish to read the original research paper from which this experiment is adapted: see Hill, N. J.; Hoover, J. M.; Stahl, S. S., *J. Chem. Educ.* 2013, 90, 102–105.

EXPERIMENTAL PROCEDURES

Oxidation of Alcohols

A * Oxidation of Cyclododecanol to Cyclododecanone



Green Experiment

Purpose To demonstrate the oxidation of a secondary alcohol to the corresponding ketone using hypochlorous acid.

SAFETY ALERT




1. Wear safety glasses or goggles and suitable protective gloves while performing the experiment.
2. Do not allow the solution of sodium hypochlorite to come in contact with your skin or eyes. If it does, flush the affected area immediately with copious amounts of water. Sodium hypochlorite will also bleach clothing.

MINISCALE PROCEDURE

Preparation Refer to the online resources to answer Pre-Lab Exercises, access videos, and read the MSDSs for the chemicals used or produced in this procedure. Review Sections 2.9, 2.11, 2.13, 2.21, 2.22, and 2.29.

Apparatus A 25-mL round-bottom flask, separatory funnel, apparatus for heating under reflux, magnetic stirring, simple distillation, and *flameless* heating.


Setting Up Place 0.5 g of cyclododecanol, 1.2 mL of acetone, and 0.4 mL of *glacial* acetic acid in the round-bottom flask. Equip the flask with a stirbar, and set up the apparatus for heating under reflux.



Oxidation Stir the mixture and warm it to approximately 45 °C; maintain this temperature within ± 5 °C throughout the course of the reaction. You may wish to monitor the temperature by suspending a thermometer through the top of the condenser using a copper wire to hold the thermometer in place. Using a Pasteur pipet, add 4.5 mL of commercial bleach (ca. 5.3% sodium hypochlorite) dropwise to the stirred mixture through the top of the condenser over a period of about 0.5 h. Upon completing the addition, stop stirring and heating the mixture so the layers may separate. Using a Pasteur pipet, remove a small portion of the aqueous layer, and place a drop or two of this solution on a dampened piece of starch/iodide test paper to determine whether sufficient hypochlorite has been added. The indicator paper immediately turns blue-black in color if sufficient bleach has been added. If this color does not develop, add an additional 0.4 mL of bleach to the reaction mixture. Stir the resulting mixture with heating for 2–3 min and repeat the test for excess hypochlorite. Add additional 0.4-mL portions of bleach until a positive test for oxidant is observed. Then stir the reaction mixture with heating for an additional 10 min and retest for hypochlorite. If the test is negative, add a final 0.4 mL of bleach. Whether this last test is positive or negative, stir the mixture with heating for 10 min more to complete the reaction.

Work-Up Allow the reaction mixture to cool to room temperature, and transfer it to a separatory funnel using a Pasteur pipet. Rinse the round-bottom flask with 5 mL of diethyl ether, and use a filter-tip pipet to transfer this wash to the separatory funnel. Shake the two-phase mixture, and separate the layers. Extract the aqueous layer with an additional 5-mL portion of diethyl ether, and add this extract to the original one. Wash the combined organic extracts with 5 mL of saturated sodium bicarbonate. Before shaking this mixture, swirl the *unstopped* funnel until the evolution of carbon dioxide ceases. Shake the mixture, venting the funnel frequently to relieve any pressure that might develop. Wash the organic solution sequentially with 5-mL portions of saturated aqueous sodium bisulfite and saturated aqueous sodium chloride. Transfer the organic solution to an Erlenmeyer flask, and dry it over several spatula-tips full of *anhydrous* sodium sulfate.* Swirl the flask occasionally for a period of 10–15 min to facilitate drying; add further small portions of *anhydrous* sodium sulfate if the solution does not become clear.

Isolation and Purification Using a filter-tip pipet, transfer the dried ethereal solution to a tared 25-mL round-bottom flask, and equip it for simple distillation. Remove the diethyl ether by simple distillation. Alternatively, use rotary evaporation or other techniques to concentrate the solution. The final traces of solvent may be removed by attaching the flask to a vacuum source and gently swirling the contents as the vacuum is applied.* The oil that is initially formed after removal of the solvents should solidify. Recrystallize the cyclododecanone from aqueous methanol.



Analysis Weigh the flask and calculate the yield of solid cyclododecanone. Determine the melting point of the product. Prepare the semicarbazone (mp 218–219 °C) or oxime (mp 131–132 °C) according to the procedures given in Section 16.2C. If necessary, recrystallize the derivatives from methanol or some other appropriate solvent. Obtain IR and ^1H NMR spectra of your starting material and product, and compare them with those of authentic samples (Figs. 16.1–16.4).

WRAPPING IT UP

Flush all aqueous solutions down the drain. Spread the sodium sulfate on a tray in the hood to evaporate residual solvent from it, and then put the used drying agent in the nonhazardous solid waste container. Place the diethyl ether recovered by distillation in the container for nonhalogenated organic solvents.

EXERCISES

1. Prove that no net oxidation occurs at the carbon atoms in the transformations of Equations 16.10 and 16.11 by comparing the oxidation numbers for carbon in the reactants and products.
2. Consider the oxidation of cyclododecanol (1) to cyclododecanone (2) using hypochlorous acid to answer the following questions.
 - a. What is the oxidation number of the carbon atom bearing the alcohol functional group in 1?
 - b. What is the oxidation number of this carbon atom in 2?
 - c. What is reduced in this reaction, and to what is it reduced?
 - d. Suggest a reagent other than bleach that would oxidize 1 into 2.
 - e. Write the chemical structure of a side-product that might be formed by the α -halogenation of 2.
3. Would it be possible to convert cyclododecanone (2) to cyclododecanol (1)? If so, what type of reaction would this be and what reagent(s) might you use?
4. Why is glacial acetic acid used in this reaction?
5. What is the function of sodium bisulfite in the procedure for isolating cyclododecanone?
6. Would cyclododecanone be expected to give a positive iodoform test (Sec. 25.7E)? Explain.

Refer to the online resources for Chapter 25 to answer Question 6.